

## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <a href="http://about.jstor.org/participate-jstor/individuals/early-journal-content">http://about.jstor.org/participate-jstor/individuals/early-journal-content</a>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

1347 July 1, 1904

## Report from Ponce.

Acting Assistant Surgeon Torres reports, June 3 and 15, through the chief quarantine officer, as follows:

Summary of transactions of service during the month of May, 1904.

Vessels inspected	12
Bills of health issued	$\overline{22}$
Passengers inspected (in transit)	446
Passengers inspected (incoming)	70
Crew inspected	3 <b>4</b> 4
Vessels in quarantine	1
Immigrants inspected.	$3\hat{5}$
Rejection	1
Passengers detained in quarantine	ô
Pieces of baggage disinfected.	ő

## Vital statistics.

Number of causes of deaths in Ponce jurisdiction (city, playa, and surrounding country) during the month of May, 1904.

Diseases of:		Accidents	1
Digestive system	36	Tuberculosis	5
Nervous system	$^{2}$	Enteric fever	1
Circulatory system	6	Convulsions	5
Respiratory system	19	Without official data	18
Malarial fever	15	·	
Hydrœmias	4	Total	157
Anæmias, inanition	22	=	
Nephritis	1	May, 1903:	
Old age	3	Deaths	151
Metritis	5	Births	114
Rhachitis	6	May, 1904:	
Congenital malformation	6	Deaths	157
Cancer	$^2$	Births	149

Foreign and insularal statistic reports of countries and cities—Yearly and monthly.

Austria—Brunn.—Month of April, 1904. Estimated population, 95,342. Total number of deaths, 252, including diphtheria 1, enteric fever 2, scarlet fever 7, and 59 from tuberculosis.

Brazil—Ceara.—Month of April, 1904. Estimated population, 50,000. Total number of deaths, 185, including 14 from enteric fever.

Canada—British Columbia—Victoria.—Month of May, 1904. Estimated population, 21,000. Total number of deaths, 17, including 2 from tuberculosis.

Cuba—Habana.—Month of January, 1904. Estimated population, 273,000. Total number of deaths, 520, including diphtheria 4, enteric fever 6, measles 3, scarlet fever 14, and 114 from tuberculosis.

Formosa.—Month of April, 1904. Estimated population, 2,797,543. Total number of deaths not reported. One death from enteric fever, and 1,036 deaths from plague reported.

France—Marseille.—Month of May, 1904. Estimated population, 491,161. Total number of deaths, 799, including diphtheria 6, enteric fever 9, measles 10, scarlet fever 1, whooping cough 1, and 84 from tuberculosis.

July 1, 1904 1348

GERMANY— Weimar.—Month of May, 1904. Census population, 30,606. Total number of deaths, 49, including diphtheria 2, and 1 from enteric fever.

Great Britain—England and Wales.—The deaths registered in 76 great towns in England and Wales during the week ended June 11, 1904, correspond to an annual rate of 14.1 per 1,000 of the aggregate population, which is estimated at 15,271,287.

London.—One thousand one hundred and eighty-seven deaths were registered during the week, including measles 62, scarlet fever 4, diphtheria 9, whooping cough 28, enteric fever 5, and diarrhea 26. The deaths from all causes correspond to an annual rate of 13.3 per 1,000. In Greater London 1,617 deaths were registered. In the "outer ring" the deaths included 2 from diphtheria, 9 from measles, and 5 from whooping cough.

Ireland.—The average annual death rate represented by the deaths registered during the week ended June 11, 1904, in the 21 principal town districts of Ireland was 18.1 per 1,000 of the population, which is estimated at 1,093,289. The lowest rate was recorded in Sligo, viz, 4.8, and the highest in Kilkenny, viz, 39.3 per 1,000. In Dublin and suburbs 143 deaths were registered, including enteric fever 1, measles 1, scarlet fever 1, whooping cough 1, and 44 from tuberculosis.

Scotland.—The deaths registered in 8 principal towns during the week ended June 11, 1904, correspond to an annual rate of 16.2 per 1,000 of the population, which is estimated at 1,726,236. The lowest rate of mortality was recorded in Greenock, viz, 11.2, and the highest in Perth, viz, 29.2 per 1,000. The aggregate number of deaths registered from all causes was 536, including diphtheria 4, measles 16, scarlet fever 3, smallpox 3, and 21 from whooping cough.

HAWAII—Honolulu.—Month of May, 1904. Census population, 39,306. Total number of deaths, 80, including enteric fever 4, plague 1, and 11 from tuberculosis.

Jamaica—Kingston.—Month of May, 1904. Estimated population, 46,542. Total number of deaths, 118, including 21 from phthisis pulmonalis.

Japan—Nagasaki.—Ten days ended May 20, 1904. Estimated population, 154,727. Total number of deaths not reported. One death from diphtheria and 6 deaths from smallpox reported.

Eleven days ended May 31, 1904. Estimated population, 154,927. Total number of deaths not reported. One death from diphtheria and 7 deaths from smallpox reported.

JAVA—Batavia.—Weeks ended April 23 and May 14, 1904. Estimated population, 160,000. Total number of deaths not reported. One death from smallpox reported.

PORTO RICO.—Month of April, 1904. Estimated population,

1349 July 1, 1904

953,243. Total number of deaths 1,668, including diphtheria 1, enteric fever 10, and 139 from tuberculosis.

URUGUAY—Montevideo.—Month of April, 1904. Estimated population, 284,107. Total number of deaths, 402, including diphtheria 1, enteric fever 21, scarlet fever 48, smallpox 1, and 46 from tuberculosis.

West Indies—St. Thomas.—Weeks ended May 13 and May 27, 1904. Estimated population, 11,012. Total number of deaths, 8 and 6, respectively. No contagious diseases.

Cholera, yellow fever, plague, and smallpox, June 25 to July 1, 1904.

[Reports received by the Surgeon-General, Public Health and Marine-Hospital Service, from United States consuls through the Department of State and from other sources.]

[For reports received from December 26, 1903, to June 24, 1904, see Public Health Reports for June 24, 1904.]

[Note.—In accordance with custom, the tables of epidemic diseases are terminated semiannually and new tables begun.]

CHOLERA.

Place.	Date.	Cases.	Deaths.	Remarks.				
China: Hongkong.	May 15-28	13	12	Two cases on ss. Ping Shan, from Saigon.				
YELLOW FEVER.								
Brazil:								
Rio de Janeiro Mexico:	May 9-22	19	5					
Tampico Veracruz	June 12–18 June 25	1 1		On ss. Habana, from Progreso.				
Peru: Rio Chiclayo	May 19	2						
.,	PL	AGUE.						
Chile:			-					
Antofagasta China:	May 19	15						
Canton	May 28 May 15-28	62	60	Spreading. Two cases on unknown steam- ships from Amoy and Shek Loong.				
Formosa: Ensuiko Kagi Kelung	May 15-28 May 15-28 May 15-28	123 85 3	80 82 3					
Pescadores Shinko	May 15–28 May 15–28	1 5						
Taihoku Tainan Toroku	May 15-28 May 15-28 May 15-28	13 335 31	13 288 18					
Hawaii: Honolulu	June 21		10					
Japan: Kobe	May 31			On S. S. Korea from Hongkong				
	may of	1		via Amoy, Shanghai, and Nagasaki.				
Peru; Callao Lima Payta.	May 27-June 4 May 22-28 May 29-June 4	4 13 11	4 7	Number of cases from Oct., 1903 to May 15, 1904, 224.				
South Africa: Cape Colony (Port Elizabeth).	May 15-21	2						